

Tylenol / Acetaminophen Dosing Chart (by weight)

Can be given every 4 hours not to exceed 5 doses in 24 hours.

Weight (pounds)	mg	Infant Drops 160 mg / 5 mL	Children's 160 mg / 5 mL	Chewable 160 mg	Tablets 325 mg	Tablets 500 mg
DO NOT USE IN CHILDREN UNDER 12 WEEKS OF AGE UNLESS APPROVED BY YOUR PEDIATRICIAN.						
6 – 11 lbs.	40 mg	1.25 mL	1.25 mL			
12 – 17 lbs.	80 mg	2.5 mL	2.5 mL			
18 – 23 lbs.	120 mg	3.75 mL	3.75 mL			
24 – 35 lbs.	160 mg	5 mL	5 mL	1 tablet		
36 – 47 lbs.	240 mg	7.5 mL	7.5 mL	1 ½ tablets		
48 – 59 lbs.	325 mg	10 mL	10 mL	2 tablets	1 tablet	
60 – 71 lbs.	400 mg	12.5 mL	12.5 mL	2 ½ tablets	1 tablet	
72 – 95 lbs.	480 mg	15 mL	15 mL	3 tablets	1 ½ tablets	1 tablet
96 + lbs.	650 mg	20 mL	20 mL	4 tablets	2 tablets	1 tablet
*** Acetaminophen also comes in 80 mg, 120 mg, and 650 mg suppositories. The rectal dose is the same as the dosage given by mouth.						

Motrin / Ibuprofen Dosing Chart (by weight)

Can be given every 6-8 hours.

Weight (pounds)	mg	Infant Drops (CONCENTRATED) 50 mg / 1.25 mL	Children's 100 mg / 5 mL	Chewable 100 mg	Jr. Tablets 100 mg	Tablets 200 mg
CHILD MUST BE 6 MONTHS OF AGE BEFORE GIVING MOTRIN / IBUPROFEN						
12 – 17 lbs.	50 mg	1.25 mL	2.5 mL			
18 – 23 lbs.	75 mg	1.875 mL	4 mL			
24 – 35 lbs.	100 mg	2.5 mL	5 mL	1 tablet		
36 – 47 lbs.	150 mg	3.75 mL	7.5 mL	1 ½ tablets		
48 – 59 lbs.	200 mg	5 mL	10 mL	2 tablets	2 tablets	1 tablet
60 – 71 lbs.	250 mg		12.5 mL	2 ½ tablets	2 ½ tablets	1 tablet
72 – 95 lbs.	300 mg		15 mL	3 tablets	3 tablets	1 ½ tablets
96 + lbs.	400 mg		20 mL	4 tablets	4 tablets	2 tablets

Conversion Chart	
2.5 mL = ½ tsp.	10 mL = 2 tsp.
3.75 mL = ¾ tsp.	12.5 mL = 2 ½ tsp.
5 mL = 1 tsp.	15 mL = 3 tsp.
7.5 mL = 1 ½ tsp.	20 mL = 4 tsp.